



**A NARRATIVE REVIEW OF TRENDS OF HUMAN IMMUNODEFICIENCY VIRUS
SERO STATUS AMONG BLOOD DONORS IN SUB-SAHARAN AFRICA**

**ADDIS ADERA GEBRU¹, TADESSE BEKELE TAFESSE^{2*}, SEMGNE GOBALEE¹,
GOSAYE DEGU BELAY³, DEMELASH ATARO⁴, MOLLA TEFERI BELEW⁵, BELAY
ALI EBRAHIM⁶, GETACHEW MEKONNEN SHEBESHI⁷ AND YONAS YIMAM⁸**

¹Department of Nursing, Faculty of Health Sciences, Woldia University, Woldia, Ethiopia

²School of Pharmacy, College of Health and Medical Sciences, Haramaya University, Harar,
Ethiopia

³Department of Disasters and Emergency Health, School of Public Health, Tehran University of
Medical Sciences, Tehran, Iran.

⁴Department Emergency Medicine, Head of Emergency Unit and Coordinator of Emergency
Unit, Hawassa University, Hawassa, Ethiopia

⁵Department of Public Health, College of Health Sciences, Mekelle University, Mekelle,
Ethiopia

⁶Department of Public Health, Faculty of Health Sciences, Samara University, Samara, Ethiopia

⁷Amhara Regional Health Office, North Wollo Zone Health Department, Harbu District, Woldia,
Ethiopia

⁸Department of Biology, Faculty of Natural & Computational Sciences, Woldia University,
Woldia, Ethiopia

***Corresponding Author: Tadesse Bekele Tafesse: School of Pharmacy, College of Health
and Medical Sciences, Haramaya University, Harar, Ethiopia**

Email: tadie2007@yahoo.com

Received 29th July 2016; Revised 29th August 2016; Accepted 2nd Sept. 2016; Available online 8th Nov. 2016

ABSTRACT

Background: The prevalence of HIV is contributing the increasing morbidity and mortality in Sub-Saharan Africa region. Blood transfusion can be a life-saving intervention. But it can also causes acute or delayed complication and may bring the risk of transfusion–transmissible